

## YUMSUK JOURNAL OF PURE AND APPLIED SCIENCES

### OPTIMIZATION OF ITACONIC ACID (IA) PRODUCTION USING ORGANIC ACID PRODUCING FUNGI ISOLATED FROM SOIL

<sup>1</sup>Sadiq B.B., <sup>3</sup>Aliyu A., <sup>1</sup>Amina A. G., <sup>1</sup>Tanko N.D., <sup>2</sup>Nasiru S., <sup>1</sup>Zulaihat L., <sup>2</sup>Bala M.

<sup>1</sup> Department of Biochemistry, Bayero University, Kano

<sup>2</sup> Department of Biochemistry, Federal University, Dutse

<sup>3</sup> Department of Biochemistry, Aliko Dangote University of Science and Technology Wudil, Kano

\*Corresponding Author e-mail: sadiqbabuga63@gmail.com

**Abstract:** Microbial production of organic acids has been incessantly escalating over conventional chemical methods due to several advantages including enantio-selectivity, high purity, less environmental pollution and cost effectiveness. Itaconic acid is used as a platform chemical for the production of various value-added chemicals such as poly-itaconic acid, resins biofuel components, ionomer cements. This research work aimed to isolate *Aspergillus terreus* producing itaconic acid from soil and optimize the maximum yield conditions. *Aspergillus terreus* was isolated, screened and identified morphologically and molecularly and for itaconic acid production. The itaconic acid produced was confirmed by bromocresol purple and measured by UV spectrophotometer in which the optical density was taken at 385nm. Itaconic acid was optimized initially by one factor at a time (OFAT) and then response surface methodology (RSM) using Central Composite Design (CCD). Incubation time, substrate concentration, pH and temperature were optimized and obtained the maximum yield of 22.601mg/ml itaconic acid at incubation time of 96hrs, substrate concentration 10% of molasses, pH of 4.0 and temperature of 34.5°C with a significant quadratic model of P- value less than 0.05. The response surface plots (3D and contour) presented that the interactions between the parameters were significant to itaconic acid production. Based on this research *Aspergillus terreus* has a good potential influence for microbial production of itaconic acid using molasses as substrate.

**Key words:** Fermentation, itaconic acid, *Aspergillus terreus*, optimization, molasses

## INTRODUCTION

Increasing environmental concern and depletion of limited resources have led to the development of microbial approaches for the production of commodity chemicals for sustainable development (Aliyu *et al.* 2022). The advances in microbiology and fermentation technology have led to the development of eco-friendly processes replacing some of the conventional chemical methods. Microbial production of organic acids has been incessantly escalating over conventional chemical methods due to several advantages including enantioselectivity, high purity, less environmental pollution and cost effectiveness (Aliyu *et al.* 2022; Mari *et al.* 2022). The increasing knowledge on metabolism and pathway regulation of industrially relevant organisms has already proven to be invaluable for generating rational strain development with primarily industrial performances. In recent times, microbial production of building block chemicals is progressively expanding its market for the production of succinic, lactic, citric, itaconic, gluconic, lacto bionic acids, etc. Organic acids are utilized directly or indirectly in a wide range of applications including food, healthcare, cosmetics, textile, solvents, and construction industries as well as for the manufacturing of biodegradable packaging materials. Organic acids with wide applications in various fields are made from living cells commercially (Rajendra *et al.* 2017).

Itaconic acid is an important component in the chemical industry. Itaconic acid (IA) and its derivatives have broad application spectrum in textile, chemicals, and pharmaceutical industries. It is used as building block for acrylic plastics, acrylic latexes, anti-scaling agents and super absorbents (Steiger *et al.* 2013). It also has other applications in food packaging, detergents, paints and coatings, pharmaceuticals, agriculture, emulsifiers, herbicides and printing chemicals. Many microorganisms, such as *Ustilago zeae*, *U. maydis*, *Candida sp.*, and *Rhodotorula sp.* have been reported to produce itaconic acid (Kawamura *et al.* 1981). However, *Aspergillus terreus* is a preferred source in commercial production of itaconic acid up to 80 g/L (Steiger *et al.* 2013). The annual market of

about 80,000 tons is currently met by fungal fermentation relying on natural *Aspergillus spp.* producers (Becker *et al.* 2015).

Itaconic acid is used as a platform chemical for the production of various value-added chemicals such as poly-itaconic acid, resins biofuel components, ionomer cements etc. Itaconic acid and its derivatives have wide applications in the textile, chemical and pharmaceutical industries. The depletion of fossil fuels and the need for sustainable development require that fermentative itaconic acid production replace petroleum-based methods of itaconic acid production. Various microorganisms have been employed in itaconic acid fermentations, with the most prolific producer being *Aspergillus terreus*. Over 80 g/L itaconic acid has been produced in fermentations using glucose. However, there is an increasing interest in the utilization of lignocellulosic materials for itaconic acid production due to the concern of food security. The current industrial applications of itaconic acid and its potential use as a drop-in or novel substitute monomer to replace petroleum-based chemicals were also extensively explored. Recent trends in itaconic acid research summarized that itaconic acid can be produced cost effectively from sustainable raw materials and have the potential to replace petro-based chemicals in various applications (El-Imam *et al.* 2014).

In contrast with citric, gluconic, and lactic acids, itaconic acid is used exclusively in non-food applications. Its primary application is in the polymer industry where it is employed as a co-monomer at a level of 1–5% for certain products. Itaconic acid is also important as an ingredient for the manufacture of synthetic fibers, coatings, adhesives, thickeners, and binders. The market volume has been estimated to be about 15,000 metric tons per year and is expected to grow if the selling price (estimated to be about US\$4 per kg) can be reduced (Willke and Vorlop 2001). To date very little research has been directed at the improvement of itaconic acid production. In contrast, there has been a larger research effort directed at lactic acid production to feed the market for biodegradable plastic.

Itaconic acid was historically produced by various chemical methods which includes Destructive distillation of citric acid, and this was the main method of producing it prior to the 1960s; Oxidation of isoprene etc. None of these (or other) processes

compete favourably with the fermentative production process (Willke and Vorlop 2001), and itaconic acid (IA) is now almost entirely produced by fermentation of sugars by *Aspergillus terreus* (Tsao 1990). The Northern Regional Research Laboratory (NRRL) of the United States Department of Agriculture (USDA) screened several wild type strains and identified *Aspergillus terreus* NRRL 1960, as the most prolific itaconic acid (IA) producing strain (Lockwood and Reeves 1945) which then went on to become the most published strain.

The initial industrial production of itaconic acid used a chemical approach that is the pyrolysis of citric acid to itaconic anhydride, followed by the hydrolysis of the anhydride which leads to the production of nonrenewable chemicals. The chemical synthesis of itaconic acid (IA) is not very effective. Thus, it is estimated that the market volume for itaconic acid (IA) is about 80,000 tons per year in 2005 and the selling price is \$2.00 per kg (Okabe *et al.*, 2009), and there is an expectation for a market demand increase if the selling price can be reduced. Global carbon emissions as a result of petroleum-based processing and products are forcing industries to look for alternative processing and production methods.

Production of itaconic acid by fermentation needs significant improvements which will increase the productivity and reduction costs of the production. Bio-based itaconic acid production promises to be an attractive alternative for the chemicals industry to replace petroleum-based chemicals synthesis. Many chemicals such as succinic acid, 1, 3-propanediol and ethanol which were hitherto made from petroleum refining are now being successfully produced from renewable biomass. The high request for bio-based materials anticipates a further growth of 60% of the itaconic acid (IA) world market, which is predicted to surpass 216 million USD in the year 2020. According to the annual forecast, market is predicted to exceed 410,000 tons of itaconic acid by the year 2020 (Choi *et al.* 2015). Itaconic Acid (IA) is an important platform chemical which has a wide range of actual and potential applications. It can be used to replace a

wide range of petroleum based chemicals, e.g. acrylic acid, which will reduce dependence on petroleum and the attendant deleterious environmental effects.

## MATERIALS AND METHOD

### Sampling of the Soil

The soil sample was collected at Bayero University Kano, old campus (Latitude of 11.9742°N and Longitude of 8.4684°E) Gwale LGA, Kano Nigeria and collected from a dark loamy soil and transferred safely to the laboratory. Exactly 1 g of the soil sample was weighted and diluted with 9ml distilled water using test tubes.

### Medium Preparation

The primary medium used for isolation of fungi was potato dextrose agar (3.9 g of agar was dissolved in 100 ml of distilled water). The medium was autoclaved at 121°C and 15 lb pressure for 15 min. The medium was poured on Petri plates in sterilized laminar flow to avoid contamination for the growth of fungi.

### Isolation of *Aspergillus terreus* from the Soil

#### Sample

From the collected soil sample (Soil dilution method: Hawaz *et al.*, 2023) diluted with 1 g soil is in 10ml of sterile distilled water. Exactly 9 ml of distilled water was taken to 5 different test tubes and 1 ml of the soil liquid suspension was serially transferred to each 9 ml distilled water containing test tubes. 1 ml of suspension from each 5 different test tubes after dilution was added to sterile Petri plates in duplicates containing sterile Potato Dextrose Agar (PDA) medium. The medium was autoclaved at 121°C and 15 lb pressure for 15 min. The plates were incubated at 28°C for 5- 7 days. The growth of separate colonies was observed. A greater number of species were isolated most of the fungus speculates heavily. Pure culture was done using Petri plates in duplicates for each colony containing fresh agar of PDA medium.

### Morphological Identification of *Aspergillus terreus* Staining Reaction

The pure fungus of *Aspergillus terreus* grown as metallic brown colonies or different colours. From the bottom, the colonies confer on the medium a dark brown color. The isolated fungi were identified to the genus level and to the species when possible on the basis of macro morphology and micro morphology (Shalini *et al.* 2014).

The macro morphological is by observing the colony

features (color, shape, size and hyphae). The colonies were examined for slow or for rapid growth, topography (flat, heaped, regularly or irregularly folded), texture (yeast like, powdery, granular, velvety or cottony), surface (pigmentation and reverse pigmentation).

The micro morphological is by a compound microscope with a digital camera using a lacto phenol cotton blue-stained slide mounted with a small portion of the mycelium (Gaddeyya *et al.* 2012). The Hyphae, macro conidia, micro conidia, chlamydospores and other special fungal structure, characteristics using suitable media, slide cultures and the most updated keys for identifications. The identified fungi confirmed with microbial expert. The fungal propagules were coloured. The coloured mycelia /spores/conidia and cytoplasm were stained by using Lacto phenol and cotton blue. Cotton blue were stained cytoplasm and results in light blue background. Lacto phenol acts as a cleaning agent. The stained specimen (*Aspergillus terreus*) was observed under the light microscope (Magnus MLXi plus) for identification and microphotograph was taken under 10X × 40X magnification.

#### **Pretreatment of Molasses**

Molasses are thick brown sweet liquid that is made from raw sugar. Cane molasses was pretreated with acid ( $H_2SO_4$ ). 2.0 N of the acid was used for molasses pretreatment. 15ml of the cane molasses was diluted upto 100ml with distilled water. Then 5ml of the acid was added and placed in a water bath at  $90 \pm 2^\circ C$  for 1 hour. After cooling at room temperature, the medium was neutralized with lime (CaO) and left to stand overnight. Two layers were formed, the upper shiny black and lower yellowish brown due to the presence of trace metals. The clear supernatants were diluted to desired sugar level (Shazia *et al.*, 2015).

#### **Determination of Optimum Growth**

##### **Conditions for Itaconic Acid (IA) Production**

Four parameters (Incubation time, Substrate concentration, pH and Temperature) were considered for optimum growth of the fungal isolates for the production of itaconic acid (IA). One factor at a time (OFAT) method of optimization was performed first before using

response surface methodology (RSM) by using Design Expert version 6.6.0 optimization software (Aliyu *et al.* 2022).

#### **One Factor at a Time (OFAT) for Itaconic Acid (IA) Yield**

For optimum incubation time determination, six Erlenmeyer flasks having 100 ml of Czapek Dox broth were prepared in duplicate, and their incubation time were adjusted at 24 hrs, 48 hrs, 72 hrs, 96 hrs, 120 hrs and 144 hrs which were then autoclaved. The Erlenmeyer flasks were properly inoculated with freshly prepared spores suspension (inoculum size) of fungal isolates and incubated under proper conditions at  $35^\circ C$ , pH of 5.0 and substrate concentration (molasses) of 3% in an incubator shaker at constant rpm (200) were used to investigate the influence of itaconic acid (IA) production, their absorbance were taken at 385nm using a UV Spectrophotometer.

For determining the optimum percentage of substrate concentration (molasses), six Erlenmeyer flasks having 100 ml of Czapek Dox broth were prepared in duplicate, and their percentage substrate concentration (molasses) were adjusted to 2%, 4%, 6%, 8%, 10% and 12% which were then autoclaved. The Erlenmeyer flasks were properly inoculated with freshly prepared spores suspension (inoculum size) of fungal isolates and incubated under proper conditions at temperature of  $35^\circ C$ , pH of 5.0 and incubation time of 120 hrs in an incubator shaker at constant rpm (200) were used to investigate the influence of itaconic acid (IA) production, their absorbance were taken at 385 nm using a UV Spectrophotometer.

For optimum pH determination, six Erlenmeyer flasks having 100 ml of Czapek Dox broth were prepared in duplicate, and their pH were adjusted to 3.0, 3.5, 4.0, 5.0, 6.0 and 7.0 which were then autoclaved. The Erlenmeyer flasks were inoculated with freshly prepared spores suspension (inoculum size) of fungal isolates and incubated under proper conditions at temperature of  $35^\circ C$ , incubation time of 120hrs and substrate concentration (molasses) of 3% in an incubator shaker at constant rpm (200) were used to investigate the influence of itaconic acid (IA) production, their absorbance were taken at 385nm using a UV Spectrophotometer.

For optimum temperature determination, six Erlenmeyer flasks having 100ml of Czapek Dox broth were prepared in duplicate, and their temperature were adjusted to  $30^\circ C$ ,  $32^\circ C$ ,  $35^\circ C$ ,  $37^\circ C$ ,



40°C and 42°C which were then autoclaved. The Erlenmeyer flasks were properly inoculated with freshly prepared spores suspension (inoculum size) of fungal isolates and incubated under proper conditions at incubation time of 120hrs, substrate concentration (molasses) of 3% and pH of 5.0 in an incubator shaker at constant rpm (200) were used to investigate the influence of itaconic acid (IA) production, their absorbance were taken at 385nm using a UV Spectrophotometer (Meena *et al.*, 2010).

### Response Surface Methodology (RSM) for Itaconic Acid (IA) Yield

In general, the change in culture conditions greatly influenced the production ability of itaconic acid (IA) synthesis. The statistical base optimization was used to study the influential effect of incubation time, substrate concentration (molasses), pH and temperature on itaconic acid (IA) yield using Central Composite Design (CCD). The optimized conditions of parameters were taken as independent variables and the itaconic acid (IA) yield was chosen as the dependent variables (Table 1), this resulted to thirty experimental runs from the software Design expert, version 6.0.6. The modeling and data analysis were performed using Design expert software, version 6.0.6 (Aliyu *et al.* 2022).

**Table 1: Factors for RSM Experimental Design.**

Indicator	Factor	Low level	High level
A	Incubation time	72	120
B	Substrate conc.	8	12
C	pH	3	5
D	Temperature	32	37

### Validation of the Second Order Polynomial Model

The second order polynomial model obtained from RSM was validated by conducting a series of experiments selected at random from the design in Table 5. The experiments were done by choosing random values of parameters within

the optimized levels as presented in Table 5 below. The experimental output was then compared to the values predicted by the second order model obtained from CCD, to estimate the fitness and goodness of the model.

### Data Analysis

The average data and standard deviations were obtained from the duplicate of experiments for each run using Microsoft Excel (Office, 2019). The standard deviation for each value was 5% analysis of variance (ANOVA) was done using Design-Expert software 6.0.6. A confidence level of 95% was used in this study. Any p-value less than 0.05 was considered significant and vice versa.

## RESULTS AND DISCUSSION

### Morphological Identification of *Aspergillus terreus*

The results obtained from the morphological (macroscopic and microscopic) analysis of isolated organisms (AT2, AT4, AT6 and AT8) have shown variation in colony colours, reverse colours, margin covering and growth level. The macroscopic and microscopic characteristics of isolates show that AT2, AT4, AT6 and AT8 are *Aspergillus niger*, *Aspergillus fumigatus*, *Penicillium crysogenum* and *Aspergillus terreus* respectively. The macroscopic and microscopic characteristics of AT8 (*Aspergillus terreus*) is presented in Table 2 and Table 3

**Table 2: Screening of isolates for the production of itaconic acid (IA).**

Isolates Code	AT2	AT4	AT6	AT8
Indicator (BCP)	+	—	—	+

**KEY:** AT2 = *Aspergillus niger*, AT4 = *Aspergillus fumigatus*, AT6 = *Penicillium crysogenum*, AT8 = *Aspergillus terreus*, BCP = Bromocresol purple.

**Table 3: Morphological Characteristics of *Aspergillus terreus***

Characteristics	<i>Aspergillus terreus</i>
Hyphae	Septate and hyaline
Conidiospore	Smooth and hyaline
Conidia heads	Compact
Vesicle	Biseriate spherical
Shape	Glucose

**Table 4: Microscopic Characteristics of *Aspergillus terreus***





Characteristics	<i>Aspergillus terreus</i>
Surface colour	Light yellow to metallic dark brown colour.
Growth	Slow.
Margin covering	Half to one-third
Reverse of the colony	Appears metallic brown

**Molecular Identification of Isolated Fungi**

The results for molecular identification of the fungal isolate are presented in Figure 4.1 and 4.2. Figure 4.1 presented gel electrophoresis of the 18S ribosomal RNA (18SrRNA) gene result for AT8 showing 600 base pairs on the DNA molecular weight ladder, the sequence for the 18S rRNA of the isolate confirmed the identity of the isolate (AT8) as *Aspergillus terreus* with 99% identity with *Aspergillus terreus* MW881456 with accession number of OP8866158 after blasting in NCBI.

**Production of Itaconic Acid (IA) using One Factor at a Time (OFAT)**

Figure 1, 2, 3, and 4 presented the optimum production of itaconic acid results at different parameters (incubation time, substrate concentration, pH, and temperature) using one factor at a time (OFAT) technique.

**Figure 1:** Effect of incubation time on itaconic acid production at constant substrate concentration (3%), pH (5) and temperature (35°C) using one factor at a time (OFAT).**Figure 2:** Effect of substrate concentration on itaconic acid production at constant incubation time (96hrs), pH (5) and temperature (35°C) using one factor at a time (OFAT).**Figure 3:** Effect of pH on itaconic acid production at constant incubation time (96hrs), substrate concentration (10%) and temperature (35°C) using one factor at a time (OFAT).**Figure 4:** Effect of temperature on itaconic acid (IA) production at constant incubation time (96hrs), substrate concentration (10%) and pH (4.0) using one factor at a time (OFAT).

### Incubation Time Optimization

Figure 1 presented the result of gradual increase in itaconic acid (IA) yield with increasing incubation time from 24hrs to 96hrs and later declined at 120hrs and 144hrs. The optimum itaconic acid (IA) yield (10.09mg/ml) was obtained at 96hrs

### Substrate Concentration Optimization

Substrate concentration percentage of molasses has effect in itaconic acid (IA) yield in which it was increased with increase in substrate concentration from 2% to 10%. The optimum itaconic acid (IA) yield (16.65mg/ml) was obtained at 10%. In effect, it later declined at 12% of the substrate concentration. Figure 2 presented the result of gradual increase of itaconic acid (IA) yield by effect of substrate concentration percentage.

### pH Optimization

Increase in itaconic acid (IA) yield was observed when pH increased from 3.0 to 4.0 and massively declined from pH 5.0 to pH 8.0. The optimum yield of itaconic acid (IA) was achieved at pH of 4.0 (17.415mg/ml). The result of itaconic acid (IA) yield with effect in pH is presented in Figure 3.

### Temperature Optimization

Increase in itaconic acid (IA) yield was observed with increase in temperature from 30°C to 35°C, the optimum itaconic acid (IA) yield (19.05mg/ml) was obtained at the temperature of 35°C and then later declined from 37°C. Figure 4 presented the effect of temperature on itaconic acid (IA) yield.

### Optimization using Response Surface Methodology (RSM)

Results obtained for actual itaconic acid (IA) concentration at different conditions of incubation time, substrate concentration, pH and temperature from response surface methodology (RSM) was presented in Table 5, the results of the responses obtained for each experimental run and the predicted responses were closer to each other.

Model F-value of 19.96 implies the model is significant. There is only a 0.01% chance that a "Model F-Value" this large could occur due to

noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case A, A<sup>2</sup>, B<sup>2</sup>, C<sup>2</sup>, D<sup>2</sup>, AC, BC, BD are significant model terms. Values greater than 0.1000 indicate the model terms are not significant. If there are many insignificant model terms (not counting those required to support hierarchy), model reduction may improve your model. The "Lack of Fit F-value" of 0.90 implies the Lack of Fit is not significant relative to the pure error. There is a 58.92% chance that a "Lack of Fit F-value" this large could occur due to noise. Regression analysis produced the following second order polynomial fit with a satisfactory coefficient of determination ( $R^2 = 0.9490$ ).

Itaconic acid concentratin= +21.91+1.53A+0.36B +0.37C +0.21D -2.30A<sup>2</sup> -1.90B<sup>2</sup> -3.08C<sup>2</sup> -0.65D<sup>2</sup> - 0.56AB+0.74AC-0.57AD+0.82BC-1.22BD+0.46CD .....Equation ii




Where A, B, C and D are incubation time, substrate concentration, pH and temperature respectively. AB, AC, AD, BC, BD, CD are the interactions, and A<sup>2</sup>, B<sup>2</sup>, C<sup>2</sup>, D<sup>2</sup> are the quadratic terms.

**Table 5: Actual and predicted itaconic acid yield at different condition of incubation time, substrate concentration, pH and temperature.**

Run	Incubation time (hrs)	Substrate concentration (ml)	pH	Temperature (degree)	Experimental value (mg/ml)	Predicted value (mg/ml)
1	96.00	6.00	4.00	34.50	14.25	11.18
2	96.00	14.00	4.00	34.50	14.54	15.03
3	120.00	8.00	5.00	37.00	16.25	13.81
4	72.00	12.00	3.00	32.00	13.02	15.41
5	96.00	10.00	6.00	34.50	11.12	7.88
6	96.00	10.00	4.00	34.50	21.35	21.91
7	72.00	12.00	5.00	32.00	14.32	13.80
8	72.00	12.00	3.00	37.00	13.55	18.36
9	120.00	12.00	3.00	32.00	16.75	14.25
10	72.00	8.00	5.00	37.00	13.15	15.81
11	120.00	12.00	5.00	32.00	17.00	12.01
12	120.00	8.00	3.00	37.00	16.99	11.32
13	96.00	10.00	4.00	29.50	19.67	12.79
14	96.00	10.00	4.00	34.50	19.85	17.32
15	72.00	8.00	3.00	32.00	11.75	13.84
16	120.00	12.00	5.00	37.00	17.25	16.12
17	96.00	10.00	4.00	34.50	21.85	19.63
18	96.00	10.00	4.00	34.50	22.55	21.76
19	72.00	12.00	5.00	37.00	13.25	13.60
20	72.00	8.00	3.00	37.00	13.75	15.02
21	96.00	10.00	2.00	34.50	8.23	8.84
22	96.00	10.00	4.00	39.50	19.12	16.34
23	120.00	8.00	5.00	32.00	14.85	18.90
24	96.00	10.00	4.00	34.50	22.45	19.73
25	120.00	8.00	3.00	32.00	13.75	16.91
26	144.00	10.00	4.00	34.50	16.12	19.91
27	96.00	10.00	4.00	34.50	21.27	21.91
28	120.00	12.00	3.00	37.00	10.33	13.91
29	48.00	10.00	4.00	34.50	9.44	12.91
30	72.00	8.00	5.00	32.00	7.00	12.91

**Table 6: Analysis of Itaconic Acid Yield from Quadratic Model Analysis of Variance.**

Source	Sum of Square	Mean Square	F- Value	P > F	
<b>Model</b>	521.85	37.28	19.96	< 0.0001	Significant
A	56.24	56.24	30.11	< 0.0001	
B	3.05	3.05	1.63	0.2205	
C	3.35	3.35	1.79	0.2008	
D	1.03	1.03	0.55	0.4685	
A2	145.60	145.60	77.95	< 0.0001	
B2	99.04	99.04	53.02	< 0.0001	
C2	260.23	260.23	139.32	< 0.0001	
D2	11.60	11.60	6.21	0.0249	
AB	5.06	5.06	2.71	0.1205	
AC	8.82	8.82	4.72	0.0462	
AD	5.22	5.22	2.80	0.1153	
BC	10.82	10.82	5.79	0.0294	
BD	23.77	23.77	12.72	0.0028	
CD	3.40	3.40	1.82	0.1970	
Lack of Fit	17.98	1.80	0.90	0.5892	Not Significant
R-Squared	0.9490	Adj R-Squared	0.9015	Pred R-Squared	0.7854

**Figure 5: Response surface plots (3D and Contour) presenting the interaction between Incubation time and pH affecting itaconic acid (IA) production.****Figure 6: Response surface plots (3D and Contour) presenting the interaction between Substrate concentration and pH affecting itaconic acid (IA) production.****Figure 7: Response surface plots (3D and Contour) presenting the interaction between Substrate concentration and Temperature affecting itaconic acid (IA) production.**



### 3D-Response Surface Plots Representing the Interaction between the Variables

Interaction among the various factors and the determination of optimum condition for maximum itaconic acid (IA) production were studied by plotting three-dimensional (3D) and contour response surface plot, as presented in Figure 5, 6 and 7.

The results obtained shown the parabola shape of the 3D plot and the circular shape of contour plots indicated the interaction between incubation time and pH was significant, keeping substrate concentration and temperature constant. Figure 5 presented the results of response surface (3D and contour) obtained for the interaction between the incubation time and pH.

The results obtained shown the parabola shape of the 3D plot and the circular shape of contour plots indicated the interaction between substrate concentration and pH was significant, keeping incubation time and temperature constant. Figure 6 presented the results of response surface (3D and contour) obtained for the interaction between the substrate concentration and pH.

The results obtained shown the parabola shape of the 3D plot and the circular shape of contour plots indicated the interaction between substrate concentration and temperature was significant, keeping incubation time and pH constant. Figure 7 presented the results of response surface (3D and contour) obtained for the interaction between the substrate concentration and temperature.

### Validation of the Second Order Polynomial Model between the Experimental and Predicted Value of Itaconic Acid (IA)

The results obtained indicated that there was very good correlation between experimental and predicted values and in turn proved the validity of the models. The observed values of itaconic acid (IA) yield were compared with the values of predicted by the second order model. Table 7 presented the result of validation runs with observed and predicted values.

**Table 7:** Validation of the Second Order Polynomial Model between Experimental and Predicted Value of Itaconic Acid (IA) yield.

Ru n	Incubation time (hrs)	Substrate conc. (%)	pH	Temp (°C)	Experimenta l value (mg/ml)	Predicted value (mg/ml)
1	103.78	10.11	4.1 1	34.36	22.481	22.204
2	96.00	10.00	4.0 0	34.50	22.531	24.000
3	96.00	10.00	4.0 0	34.50	22.601	22.550

### DISCUSSION

In this research study, four different fungal isolates were isolated from soil and identified morphologically and molecularly. AT8 (*Aspergillus terreus*) show the highest itaconic acid production capability among the isolates (Helia and Wan 2015) and (Meena *et al.* 2010). Molasses is used as source of glucose which is a very convenient raw material for itaconic acid production (Lockwood and Reeves 1945). DNA sequencing and molecular identification from Genbank identified isolate AT8 as new strain of *Aspergillus terreus* with closet similarity of 98% identity with *Aspergillus terreus* MW881456. *Aspergillus terreus* assigned the accession number of OP866152. Sequence alignment and evolutionary history was gathered for plotting phylogenetic tree.

Itaconic acid was produced under the influence of physicochemical parameters which were determined to have high yield by optimizing them with conventional technique, one factor at a time (OFAT) and response surface methodology (RSM) (Sadiq *et al.* 2017). The parameters were maintained significantly for better fermentation yield.

Effect of incubation time on itaconic acid (IA) was observed by varying time from 24hrs to 144hrs and keeping all other variables constant. The optimum yield of itaconic acid (IA) was determined at 96hrs of incubation and declined with increase incubation time as presented in Figure 4.3, this was in line with the study done by Linda (2021) by determining the optimum incubation time of the fungal isolates. The itaconic acid yield declined due to the relationship between the *Aspergillus terreus* and sugar contents in the medium and the fungal growth curve. Microbial production of metabolites usually starts after a lag

phase of one day and reaches maximum at the onset of stationary phase or late.

Effect of substrate concentration on itaconic acid (IA) was observed and the optimum itaconic acid produced (16.65mg/ml) was obtained at 10% concentration of molasses. The *Aspergillus terreus* cell number increased exponentially which could provide the maximal conversion of substrate to itaconic acid was found at 10% this was related with the work of Meena *et al.* (2010). Low itaconic acid (IA) produced at molasses concentration above 10% and this may occur due to the formation of secondary byproducts that limit itaconic acid (IA) productivity in accordance to the research done by Hawaz *et al.* (2023).

Effect of medium pH was revealed to have the optimum itaconic acid (IA) yield of 17.415mg/ml at pH of 4.0, which was found to be similar with the work of EL Imam *et al.* (2013) and Sudarkodi *et al.* (2012) who reported a maximum yield of itaconic acid (IA) found to be at pH 4. The enzymatic reactions in the utilization of energy are regulated by pH. Basically, the impact of low pH is associated with the activity of enzymes taking part in the biosynthesis of itaconic acid (IA) and subsequent transfer mechanism to the extracellular space/out of the cell. This is related to the finding of Peter *et al.* (2019) reported that itaconic acid (IA) generation by filamentous fungi such as *Aspergillus terreus* favors lower pH conditions and it has been argued that besides enabling the appropriate growth of *Aspergillus terreus*, such a fermentation environment can be useful for suppressing the formation of by-products that would lower the final itaconic acid (IA) yield and productivity.

Effect of temperature on itaconic acid (IA) was also observed and the maximum itaconic acid (IA) production of 19.05mg/ml was obtained at 35°C. After the optimum temperature the overall growth rate began to fall due to increase in rate of microbial death, as the death rate is also a function of temperature as reported by Meena *et al.* (2010). This high value of cell death increases with increase in temperature, than the

growth rate. Hence the overall growth rates rapidly declined above the optimal temperature. Apart from this, the product inhibition effect is also more at higher temperatures than at lower temperatures.

The results obtained from the preliminary optimization using one factor at a time (OFAT) experiment were then applied to response surface methodology (RSM) modelling. 30 experimental runs using central composite design (CCD) were design as presented in Table 5, the experimental yield was recorded and relatively closed to the predicted value presented in Table 5. The statistical analysis for significances of all factors was described by analysis of variance (ANOVA) in Table 6. Based on the result obtained, the model of analysis was confirmed significant and highly reliable at P-value ( $< 0.0001$ ) less than 0.05, and the no significant of lack of fitness indicated that the model was excellent fitted with no significant noise, the  $R^2$  and adjusted  $R^2$  value were all closed to 1 (0.9490 and 9015 respectively) showing the goodness of the model (Table 6). Additionally, the significant of the interaction between incubation time and pH (AC), substrate concentration and pH (BC) and that of substrate concentration and temperature were presented in ANOVA result (Table 6) with P-value 0.0462, 0.0294 and 0.0028 respectively. Furthermore, the response surface plots (3D and contour) reveal that the interaction between the factors (incubation time, substrate concentration, pH and temperature) were all significant to itaconic acid (IA) production as presented in Figure 5, 6, and 7. This shown that the itaconic acid (IA) production was dependent on the four parameters optimized.

Validation of the second order polynomial model confirmed the optimum itaconic acid (IA) yield of (22.601mg/ml) and indicated linear interaction and quadratic effect of variables on itaconic acid production. Therefore, the developed model is considered reliable.

The experimental and the predicted value were relatively closed indicating the relative fitness of experimental model. Moreover, this shows that extraneous factor terms in a derived model equation will affect in some reduction in the calculation of the error sum of squares (Mohamed *et al.* 2013).

## CONCLUSION

According to the results obtained from this study, the isolated fungal species (*Aspergillus terreus*) has the potential ability of utilizing molasses as substrate for itaconic acid (IA) production. The study revealed that itaconic acid (IA) could be effectively yielded by adjusting conditional parameters, such as temperature, pH, incubation duration, and substrate concentration.

## REFERENCES

- Aberkane, M., Cuenca-Estrella A., Gomez-Lopez E., Petrikkou E., Mellado A., Monzón J. and L. Rodriguez. (2002). Comparative evaluation of two different methods of inoculum preparation for antifungal susceptibility testing of filamentous fungi. *Journal of Antimicrobial Chemotherapy* 50, 719–722.
- Ajiboye, A. E., Adedayo, M. R., Babatunde, S. K., Odaibo, D. A., Ajuwon, I. B., & Ekanem, H. I. (2018). Itaconic acid production from date palm (*Phoenix dactylifera* L) using fungi in solid state fermentation. *Journal of the College of Pure and Applied Sciences*, 6(1), 20-35.
- Aliyu A, Nasiru S, J U Mari, B B Sadiq, ND Tanko, Ibrahim HM, M Bala (2022). Optimization of Valine Production Using *Bacillus Cereus* Isolated from Soil. *International Journal of Biochemistry*, 5(2): 37-49. DOI: 10.36348/sijb.2022.v05i02.002
- Alonso, S., Rendueles M., and Díaz M. (2015). Microbial production of specialty organic acids from renewable and waste materials. *Critical Reviews in Biotechnology*. 34 (4), 497–513.
- Bafana, R., and Pandey R.A. (2018). New approaches for itaconic acid production: bottlenecks and possible remedies. *Critical Reviews in Biotechnology*, 38(1): 68–82. DOI: 10.1080/07388551.2017.1312268.
- Batti, M., and Schweiger, L B. (1963). Process for the production of itaconic acid. US Patent, 3, 078,217.
- Baup, S. Uebereineue Pyrogen-Citronensäure, und über Benennung der Pyrogen-Säuren überhaupt. (1837). *Annals of Chemistry and Physics*, 39–41.
- Becker, J., Lange, A., Fabarius, J., & Wittmann, C. (2015). Top value platform chemicals: bio-based production of organic acids. *Current Opinion in Biotechnology* 36:168-175.
- Blazeck, J, Hill A, Jamoussi M, et al., (2015). Metabolic engineering of *Yarrowia lipolytica* for itaconic acid production. *Metabolic Engineering*, 32: 66–73. DOI: 10.1016/j.ymben.2015.09.005.
- Choi, S., Song, C. W., Shin, J. H., & Lee, S. Y. (2015). Biorefineries for the production of top building block chemicals and their derivatives. *Metabolic Engineering*, 28, 223–239.
- Da Cruz, J.C., Camporese Sérvulo, E.F. and Castro, A.M. (2017). Microbial Production of Itaconic Acid. In *Microbial Production of Food Ingredients and Additives*; Elsevier: Amsterdam, the Netherlands, pp. 291–316, ISBN 9780128115206.
- De Pretto, C., Giordano, R.L.C., Tardioli, P.W. and Costa, C.B.B. (2018). Possibilities for producing energy, fuels, and chemicals from soybean: a biorefinery concept. *Waste Biomass Valorization* 9, 1703–1730.
- Delidovich I, Hausoul P.J.C, and Deng L. (2016). Alternative monomers based on lignocellulose and their use for polymer production. *Chemical Reviews*, 116(3): 1540–1599. DOI: 10.1021/acs.chemrev.5b00354.
- Dwiarti L., Otsuka M., Miura S., Yaguchi M., and Okabe M. (2007). Itaconic acid production using sago starch hydrolysate by *Aspergillus terreus* TN484-M1. *Bioresources and Technology*, 98, 3329–3337.
- Eimhjellen, K.E. and Larsen H. (1955). The mechanism of itaconic acid formation by *Aspergillus terreus*. 2. The effect of substrates and inhibitors. *Biochemistry Journal*, 60, 139–147
- El-Imam, A.A. and Du C. (2014). Fermentative itaconic acid production. *Journal of Biodiversity and Bioprospect. Development*, 1, 119 <https://doi.org/10.4172/ijbbd.1000119>.

- Fawcett, H.H. and Kirk-Othmer. (1985). Concise encyclopedia of chemical technology. *Journal of Hazardous Materials*, 24–25.
- Galgiani, J. N. and D. A. Stevens. (1978). Turbidimetric studies of growth inhibition of yeasts with three drugs: inquiry into inoculum-dependent susceptibility testing, time of onset of drug effect, and implications for current and newer methods. *Antimicrobial Agents Chemother* 13:249-254.
- Gonzalez-Garcia, R.-A., McCubbin T., Navone L. Stowers C. Nielsen L.-K. and Marcellin, E. (2017). Microbial propionic acid production. *Fermentation*, 3, 21.
- Kautola, H. Vahvaseleka, M., Linko. Y.Y. and P. Linko. (1985). *Biotechnology Letters*, 7, 167-1728
- Hawaz E., Mesfn Tafesse, Anteneh Tesfaye, Solomon Kiros, Dereje Beyene, Gessesse Kebede, Teun Boekhout, Marizeth Groenwald, Bart Theelen, Ayantu Degefe, Sisay Degu, Alene Admasu, Biru Hunde and Diriba Muleta. (2023). Optimization of bioethanol production from sugarcane molasses by the response surface methodology using *Meyerozyma caribbica* isolate MJTm3. Institute of Biotechnology, Addis Ababa University, Addis Ababa, Ethiopia. *Annals of Microbiology*.
- Hegde, K., Prabhu, A., Sarma S.J., Brar S.K. and Venkata Dasu V. (2016). *Potential Applications of Renewable Itaconic Acid for the Synthesis of 3-Methyltetrahydrofuran*; Elsevier Inc. Amsterdam. The Netherlands, ISBN 9780128029800.
- Helia, H., and Wan Mohtar Wan Yusoff. (2015). School of Bioscience and Biotechnology, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia. *Current Research Journal of Biological Sciences* 7(2): 37-42.
- Hevekerl, A, Kuenz A. and Vorlop K, 2014. Influence of the pH on the itaconic acid production with *Aspergillus terreus*. *Applied Microbiology and Biotechnology*, 98(24): 10005–10012. DOI: 10.1007/s00253-014-6047-2.
- Hevekerl, A., Kuenz A., and Vorlop K. D. (2014). Influence of the pH on the itaconic acid production with *Aspergillus terreus*. *Applied Microbiology and Biotechnology*, 98, 10005–10012.
- Kane, J.H., Finlay, A.C. and Amann, P.F. (1945). Production of itaconic acid, *US-Patent*, 2 385 283
- Mari, J.U., Aliyu, A., Nasiru, S., Muhammad, A.B., Ibrahim, A.A., Magaji, H. and Bala, M. (2022). Methionine Production and Optimization Using *Bacillus cereus* Isolated From Soil. *Sch Int J Biochem*, 5(7): 95-102. DOI: 10.36348/sijb.2022.v05i07.001
- Jang, Y. S., Kim B., Shin J. H., Choi Y. J., Choi S., Song C. W., Lee J., Park H.G. and Lee S.Y. (2012). Bio-based production of C2–C6 platform chemicals. *Biotechnology and Bioengineering*, 109, 2437–2459.
- Yang J., Hao Xu, Jianchun Jiang, et al., (2019). Production of itaconic acid through microbiological fermentation of inexpensive materials. *Journal of Bioresources and Bioproducts*, 4(3): 135–142. DOI: 10.12162/jbb.v4i3.001.
- Karaffa, L.; Díaz, R.; Papp, B.; Fekete, E.; Sándor, B.; Kubicek, C. P. (2015). A deficiency of manganese ions in the presence of high sugar concentrations is the critical parameter for achieving high yields of itaconic acid by *Aspergillus terreus*. *Applied Microbiology and Biotechnology*. 99, 7937–7944.
- Kinoshita, K. and Über die. (1932). Production von Itaconsäure und Mannit durch einen neuen Schimmelpilz, *Aspergillus itaconicus*. *Acta Phytchim.*, 5, 271–287.
- Klement, T., Milker, S. and Jager, G. (2012). Biomass pretreatment affects *Ustilago maydis* in producing itaconic acid. *Microbial Cell Factories*, 11(1): 43. DOI: 10.1186/1475-2859-11-43.
- Kuenz, A., Gallenmüller, Y. and Willke, T. (2012).



- Microbial production of itaconic acid: developing a stable platform for high product concentrations. *Applied Microbiology and Biotechnology*, 96(5): 1209–1216. DOI: 10.1007/s00253-012-4221-y.
- Kuenz, A. and Krull S. (2018). Biotechnological production of itaconic acid: things you have to know. *Applied Microbiology and Biotechnology*, 102(9): 3901–3914. DOI: 10.1007/s00253-018-88957.
- Kumar, S., Stecher G. and Tamura K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33, 1870–1874.
- Lockwood, L.B. and Reeves, M.D. (1945). *Arch. Biochem.*, 6, 455–469
- Lai, L. S. T., Hung C. S. and Lo C. C. (2007). Effects of lactose and glucose on production of itaconic acid and lovastatin by *Aspergillus terreus* ATCC 20542. *Journal of Bioscience and Bioengineering*, **104**, 9–13.
- Larsen, H. and Eimhjellen, K.E.K. (1955). The mechanism of itaconic acid formation by *Aspergillus terreus*. 1. The effect of acidity. *Biochemistry Journal*, 6217, 135–139.
- Li, A. Pfler, N. and Zuijderwijk, R. (2013). Reduced by-product formation and modified oxygen availability improve itaconic acid production in *Aspergillus niger*. *Applied Microbiology and Biotechnology*, 97(9): 3901–3911. DOI: 10.1007/s00253-012-4684-x.
- Li, A., Pfler N., Zuijderwijk R. and Punt, P. (2012). Enhanced itaconic acid production in *Aspergillus niger* using genetic modification and medium optimization. *BMC Biotechnology*, 12, 57.
- Lin, Y.H., Li, Y. and Huang, M.C. (2004). Intracellular expression of Vitreoscilla hemoglobin in *Aspergillus terreus* to alleviate the effect of a short break in aeration during culture. *Biotechnology Letters*, 26(13): 1067–1072. DOI: 10.1023/b:bile.0000032964.15178.7c.
- Lin, Y.H., Li, Y.F., Huang, M.C. and Tsai, Y.C. (2004). Intracellular expression of Vitreoscilla hemoglobin in *Aspergillus terreus* to alleviate the effect of a short break in aeration during culture. *Biotechnology Letters*, 26: 1067–1072.
- Linda, B. (2021). College of agriculture and college of science, department of microbiology, Origen state university. *Microbial growth and Microbiology*, LibreTexts.
- Mattey, M. (1992). The production of organic acids. *Critical Reviews in Biotechnology*, 12(1/2): 87–132. DOI: 10.3109/07388559209069189.
- Steiger, M.G., Blumhoff, M.L., Mattanovich, D. and Sauer, M. (2015). Biochemistry of microbial itaconic acid production. Austrian Centre of Industrial Biotechnology (ACIB GmbH), Department of Biotechnology, BOKU – Vienna. *Institute of BioTechnology*, University of Natural Resources and Life Sciences, Vienna, Austria.
- Meena V., Sumanjali A., Dwarka K., Subburathinam K. M. and Sambasiva Rao K.R.S. (2010). Production of itaconic acid through submerged fermentation employing different species of *Aspergillus*, 3, 100–109.
- Mondala, A. H. (2015). Direct fungal fermentation of lignocellulosic biomass into itaconic, fumaric, and malic acids: Current and future prospects. *Journal of Industrial Microbiology and Biotechnology*, 42, 487–506.
- Murali, N., Srinivas K. and Ahring B. K. (2017). Biochemical production and separation of carboxylic acids for biorefinery applications. *Fermentation*, 3, 22.
- Nelson, G.E.N. Trauffer D. H., Kelley S. E. and Lockwood L. B. (2005). Production of Itaconic Acid by *Aspergillus terreus* in 20-Liter Fermentors. *Industrial Engineering and Chemistry*, 44, 1166–1168.
- Nghiem, N. P., Kleff S. and Schwegmann S. (2017). Succinic acid: *Technology development and commercialization*. *Fermentation*, 3, 26.
- Nubel, R.C. and Ratajak, E.J. (1962). Process for producing itaconic acid. *US Patent*, 3, 044,941.

- Papagianni, M. (2007). Advances in citric acid fermentation by *Aspergillus niger*: biochemical aspects, membrane transport and modeling. *Biotechnology Advances*, 25(3): 244–263. DOI: 10.1016/j.biotechadv.2007.01.002.
- Pedroso, G.B., Montipó S., Mario D.A.N., Alves S.H. and Martins A. F. (2017). Building block itaconic acid from left-over biomass. *Biomass Convers. Biorefnery*, 7, 23–35.
- Raghu, C. and Raghuveer P. (2017). Itaconic acid Production. A short review. *International Journal of Advanced Engineering Technology Management and Applied Science*, 4, 8–15.
- Rajendra, S., Anshumali, M. Manoj, K. and Praveen Kumar Mehta (2017). Organic acids: An overview on microbial production. *International Journal of Advanced Biotechnology and Research (IJBR)* ISSN 0976-2612, Online ISSN 2278–599X, Vol-8, Issue-1, 2017, 104–111 <http://www.bipublication.com>
- Saha, B. C. (2017). Emerging biotechnologies for production of itaconic acid and its applications as a platform chemical. *Journal of Industrial Microbiology and Biotechnology*, 44, 303–315.
- Saitou, N. and Nei M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4, 406–425.
- Sarker, T.C., Azam S. M., G. G. and Bonanomi G. (2017). Recent advances in sugarcane industry solid by-products valorization. *Waste Biomass Valorization* 8, 241–266.
- Sauer, M., Porro D., Mattanovich D. and Branduard, P. (2008). Microbial production of organic acids: expanding the markets. *Trends Biotechnology* 26 (2), 100–108.
- Shalini, R. V. and K. Amutha. (2014). Identification and Molecular Characterization of *Aspergillus fumigatus* from Soil. *Journal of Medical and Pharmaceutical Innovation*; 1 (4) 2014; 12–15.
- Shazia, A., Sikander A., and Ikram-ul-Haq, A. (2015). Acidic pre-treatment of sugarcane molasses for citric acid production by *Aspergillus niger* NG-4. *International journal of current microbiology and applied sciences*.
- Shin, W.S., Lee, D., Kim, S. Jeong, Y.S. and Chun, G.T. (2013). Application of scale-up criterion of constant oxygen mass transfer coefficient (kLa) for production of itaconic acid in a 50 L pilot-scale fermentor by fungal cells of *Aspergillus terreus*. *Journal of Microbiology and Biotechnology*, 23, 1445–1453
- Sivagnanam, S. and Jayanthi, A. (2013). Ecofriendly Method for Bioremediation of Chlorpyrifos from Agricultural Soil by Novel Fungus *Aspergillus terreus* JAS1. *Microbial Biotechnology Laboratory, School of Biosciences and Technology, Division of Environmental Biotechnology, VIT University, Vellore 632014 Tamil Nadu, India.*
- Steiger, M.G., Blumhoff, M.L., Mattanovich, D. and Sauer, M. (2013). Biochemistry of microbial itaconic acid production. *Frontiers in microbiology* 4:23.
- Sudarkodi, C., Subha K., Kanimozh. K. and Panneerselvam. (2012). Optimization and production of itaconic acid using *Aspergillus flavus*. *A PG and Research Department of Botany and Microbiology*, A. V. V. M Sri Pushpam College, Poondi, Thanjavur, Tamilnadu India.
- Tamura, K., Nei, M., and Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences (USA)*, 101, 11030–11035.
- Tate, B.E. (1967). Polymerization of itaconic acid and derivatives BT-Fortschritte der Hochpolymeren-Forschung. *Advanced Polymerization in Science*, 5, 214–232.
- Tsao, G.T. (1999) Recent Progress in Bioconversion

- of Lignocellulosics. *Advances in Biochemical Engineering Biotechnology*: 65.
- Vassilev, N, Kautola H. and Linko Y, (1992). Immobilized *Aspergillus terreus* in itaconic acid production from glucose. *Biotechnology Letters*, 14(3): 201–206.
- Wasewar, K.L., Shende, D. and Keshav, A., (2011). Reactive extraction of itaconic acid using tri-n-butyl phosphate and aliquat 336 in sunflower oil as a non-toxic diluent. *Journal of Chemical Technology and Biotechnology*. 86 (2), 319–323.
- West, T.P. (2017). Microbial production of malic acid from biofuel-related coproducts and biomass. *Fermentation*, 3, 14.
- Willke, T, and Vorlop, K.D. (2001). Biotechnological production of itaconic acid. *Applied Microbiology Biotechnology*, 56: 289-295.
- Willke, T. and Kuenz, A. (2007). Itaconsäureherstellung aus nachwachsenden Rohstoffen als Ersatz für petrochemisch hergestellte Acrylsäure. Bundesforschungsanstalt für Landwirtschaft, FAL.
- Willke, T. and Vorlop, K.D. (2001). Biotechnological production of itaconic acid. *Applied Microbiology and Biotechnology*, 56(3-4), 289–295.
- Yang, L., Lübeck M. and Lübeck P. S. (2017). *Aspergillus* as a versatile cell factory for organic acid production. *Fungal Biology Review*, 31, 33–49.
- Yang, W., Hu Y., Chen Z., Jiang X., Wang J. and Wang R. (2012). Solubility of itaconic acid in different organic solvents: Experimental measurement and thermodynamic modeling. *Fluid Phase Equilibrium*, 314, 180–184.
- Zhang, K., Zhang B. and Yang S.T. (2013). Production of citric, itaconic, fumaric, and malic acids in filamentous fungal fermentation. In: Yang ST, El-Enshasy HA, Thongchul N (eds) *Bioprocessing technologies in biorefinery for sustainable production of fuels, chemicals and polymers*. John Wiley & Sons Inc, Hoboken, pp 375–397.
- Zhao, M.L., Lu X.Y. and Zong H. (2018). Itaconic acid production in microorganisms. *Biotechnology Letters*, 40(3): 455– 464. DOI: 10.1007/s10529-017-2500-5. *Performance of CdS / CdTe junctions on ZnO nanorod arrays*. 176(November 2017), 100–108.