

YUMSUK

JOURNAL OF PURE AND APPLIED SCIENCES

NEW EXACT SOLUTIONS FOR TIME-BETA FRACTIONAL BOUSSINESQ-LIKE EQUATIONS, USING THE SINE-COSINE METHOD

¹Balili, A.

¹Mathematics Department, Yusuf Maitama Sule University, Kano

*Corresponding Author e-mail: adobalili@gmail.com

ABSTRACT

The major goal of this paper is to extract exact travelling wave solutions and investigate the effects the fractional parameter on the dynamic response of soliton waves of four-time beta fractional Boussinesq-like equations. Sine-cosine method has been used to achieve explicit soliton solutions of these equations that emerged in coastal and ocean engineering. The obtained solutions have been studied in the form of hyperbolic and trigonometric functions. The behavior of some of the soliton solutions are demonstrated via 2D and 3D graphs. As a result of the fractional effects, physical changes are observed. The obtained results show that the proposed method is more convenient, powerful and efficient than other analytical approaches. The extracted results might improve our understanding of how waves propagate and could be beneficial to coastal and ocean engineering as well as other fields.

Keywords: Sine-cosine method; The Boussinesq-like equations; Time beta fractional derivative; the exact solutions.

1. INTRODUCTION

Mostly natural phenomena occurring in the universe are interpreted by nonlinear differential equations of integer order as well as non-integer order. Nonlinear fractional differential equations (NFDEs) respond promptly and effectively in a variety scientific and engineering fields, including dynamical systems, electromagnetic, technology, fusion plasma, viscoelastic, biology, signal processing, electrochemical, optical fiber, oceanography, solid state physics, geochemistry, finance, among others. Many intriguing aspects of fractional calculus and flexibility of fractional theory have captivated the attention of many researchers (Sadiya *et al.* 2022; Khantun *et al.*, 2022).

Fractional derivatives are defined in several ways, including conformable derivatives (Khalil *et al.*, 2014), Atangana-Baleanu derivatives (Atangana and Baleanu, 2016), beta derivatives (Yepez-Martinez *et al.*, 2018; Atangana *et al.*, 2016), M-truncated derivative (Sousa *et al.*,

2018), and Modified RL fractional derivatives (Jumarie, 2009). A fractional derivative can have a variety of properties, so that it is possible to use the one that is most suitable for the problem at hand. In fact, the study of fractional derivative operators is a popular topic. A lot of research has been done in this area, leading to an excessive number of findings e.g. see in (Ghanbari and Baleanu, 2020; Khater and Ghanbari, 2021; Ghanbari, 2019; Ghanbari *et al.*, 2019).

The areas that focus on the examination of wave patterns in the physical world are fascinating since they address an extremely scientific and developed idea of soliton waves. The soliton wave is an important nonlinear phenomenon. Solitons offer an intriguing perspective on nonlinear physical processes. Finding solitons of nonlinear phenomena has recently gained popularity among mathematicians and scientists. The primary advantage of soliton solutions is the fact that they can be used for evaluation equations with various kind of nonlinearities that are both integrable and non-integrable. The exact soliton solutions of fractional differential equations (FDEs) has been crucial in many kinds of physical science research. Consequently,

several approaches have been devised for computing FDEs solutions. Among them are: the extended tanh-function method (Zaman *et al.*, 2022a, b), generalized $(\frac{G'}{G})$ -expansion method (Uddin *et al.*, 2022), modified F- expansion method (Ahmad *et al.*, 2023), Sardar sub equation method (Ali *et al.*, 2023), improved F-expansion method (Akram *et al.*, 2023) extended simple equation method (Ahmad *et al.*, 2023), unified method (Ali *et al.*, 2023) and so on.

In this paper, we utilize the Sine-cosine technique introduced by Wazwaz (2004) to derive a set of new exact solutions for four distinct types of time beta fractional Boussinesq-like equations. For further context, relevant works include those by (Wazwaz, 2012), Elsami *et al.* (2014), Darvishi *et al.* (2017), Darvishi *et al.* (2018) and Osman (2019), each contributing to the understanding and development of this approach. The equations under consideration are expressed in the following forms:

$$D_{tt}^{2\beta} u - u_{xx} - (6u^2 u_x + u_{xxx})_x = 0, \quad (1)$$

$$D_{tt}^{2\beta} u - u_{xx} - (6u^2 u_x + D_{tt}^{2\beta} u_x)_x = 0, \quad (2)$$

$$D_{tt}^{2\beta} u - D_t^\beta u_x - (6u^2 u_x + D_t^\beta u_{xx})_x = 0, \quad (3)$$

$$D_{tt}^{2\beta} u - (6u^2 u_x + u_{xxx})_x = 0. \quad (4)$$

Where $D_t^\beta u$ is the time β - fractional derivative of u (Atangana's conformable) order $0 < \beta < 1$ in time $t > 0$, $u_x = \frac{\partial u}{\partial x}$, and $u_{xx} = \frac{\partial^2 u}{\partial x^2}$.

Eq. (1) includes the fourth spatial derivative u_{xxxx} and the dissipative term u_{xx} . In Eq. (2), u_{xxxx} is replaced by a mixed spatial-temporal term $D_{tt}^{2\beta} u_{xx}$. In Eq. (3), u_{xx} is replaced by $D_t^\beta u_x$ and u_{xxxx} is replaced by a spatial-temporal term $D_t^\beta u_{xx}$. Lastly, Eq. (4) lacks the dissipative term u_{xx} .

These equations (1)-(4) are non-integrable and are used as models in ocean and coastal sciences when $\beta = 1$. Some of these equations are employed in wave modeling and mathematical modeling of tidal oscillations.

This paper is structured as follows: A brief definition and properties of beta fractional derivative are given in section 2. Section 3 provides a detailed explanation of the Sine-cosine method. Section 4 discusses the application of the Sine-cosine method to the time β - fractional Boussinesq-like equations represented by equations (1)-(4) respectively. Section 5 offers a physical interpretation of some of the solutions obtained. Finally, Section 6 summarizes the main conclusions drawn from the study.

2. PROPERTIES OF BETA DERIVATIVE

Some properties of the Beta derivative are given using the following definition and theorem:

Definition (2.1), (Atangana *et al.*, 2016):

$$D_t^\beta (u(t)) = \lim_{\varepsilon \rightarrow 0} \frac{u\left(t + \varepsilon \left(t + \frac{1}{\Gamma(\beta)}\right)^{1-\beta}\right) - u(t)}{\varepsilon},$$

Where $\Gamma(\beta) = \int_0^\infty t^{\beta-1} e^{-t} dt$, $0 < \beta < 1$.

and

$$D_t^\beta (u(t)) = \frac{d^\beta u(t)}{dt^\beta}.$$

Theorem: (2.1) Let $u(t)$ and $v(t)$ be β - differentiable functions, for all $\beta \in (0,1)$ and $t > 0$. Then

$$(a) D_t^\beta (au(t) + bv(t)) = aD_t^\beta u(t) +$$

$$bD_t^\beta v(t), \forall a, b \in \mathbb{R},$$

$$(b) D_t^\beta (u(t) \cdot v(t)) =$$

$$u(t)D_t^\beta (v(t)) + v(t)D_t^\beta (u(t)),$$

$$(c) D_t^\beta \left(\frac{u(t)}{v(t)}\right) =$$

$$\frac{v(t)D_t^\beta (u(t)) - u(t)D_t^\beta (v(t))}{v(t)^2},$$

$$(d) D_t^\beta (k) = 0, \text{ where } k \in$$

$$\mathbb{R},$$

$$(e) D_t^\beta (u(t)) = \left(t + \frac{1}{\Gamma(\beta)}\right)^{1-\beta} \frac{du(t)}{dt}.$$

Relevant works on this context of beta fractional derivative see (Yusuf, *et al.*, 2019; Yiasir Arafat, *et al.*, 2023; Fiza *et al.*,

2024).

3. METHODOLOGY

In this section, we give a description of the sine-cosine method which is used to obtain exact solutions of partial differential equations.

3.1 The Sine-Cosine Method

Consider a nonlinear partial differential equation

$$P(u, u_t, u_x, u_{xx}, u_{xt}, u_{tt}, \dots D_t^\beta u, D_x^\beta u, D_{tt}^{2\beta} u, D_{xx}^{2\beta} u, \dots) = 0, 0 < \beta < 1. \quad (5)$$

Which describes the dynamical wave solution $u(x, t)$. The steps of the sine-cosine method has been proposed in (Wazwaz, 2004) as follows:

Step 1: To find the travelling wave solution of equation (5), we introduce

$$u(x, t) = U(z), \text{ and } z = \varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta. \quad (6)$$

Step 2: we use the following changes

$$\frac{\partial}{\partial t} = -\omega \frac{d}{dz}, \quad \frac{\partial^2}{\partial t^2} = \omega^2 \frac{d^2}{dz^2}, \dots, \quad \frac{\partial}{\partial x} = \varpi \frac{d}{dz}, \quad \frac{\partial^2}{\partial x^2} = \varpi^2 \frac{d^2}{dz^2} \dots \quad (7)$$

Now, utilizing Eq. (7) transforms the partial differential equation Eq. (5) into an ordinary differential equation:

$$Q(U, U', U'', \dots) = 0, \quad (8)$$

Where U' denotes $\frac{du}{dz}$.

Step 3: Simplify Eq. (8) by integration if possible

Step 4: The solution will be expressed in the following form:

$$u(x, t) = \lambda \sin^\mu(\vartheta z), \quad |z| \leq \frac{\pi}{\vartheta}, \quad (9)$$

or in the form

$$u(x, t) = \lambda \cos^\mu(\vartheta z), \quad |z| \leq \frac{\pi}{2\vartheta}, \quad (10)$$

Where λ, ϑ , and μ are parameters to be determined.

Step 5: Hence, the derivatives of Eq. (9) take the following form:

$$U(z) = \lambda \sin^\mu(\vartheta z), \quad (11)$$

$$U^n(z) = \lambda^n \sin^{n\mu}(\vartheta z), \quad (12)$$

$$U_z^n(z) = n\mu\vartheta\lambda^n \cos(\vartheta z) \sin^{n\mu-1}(\vartheta z), \quad (13)$$

$$U_{zz}^n(z) = -n^2\vartheta^2\mu^2\lambda^2 \sin^{n\mu}(\vartheta z) + n\vartheta^2\lambda^2\mu(n\mu - 1)\sin^{n\mu-2}(\vartheta z). \quad (14)$$

And the derivative of Eq. (10) becomes

$$U(z) = \lambda \cos^\mu(\vartheta z), \quad (15)$$

$$U^n(z) = \lambda^n \cos^{n\mu}(\vartheta z), \quad (16)$$

$$U_{zz}^n(z) = -n^2\vartheta^2\mu^2\lambda^2 \cos^{n\mu-1}(\vartheta z) + n\vartheta^2\lambda^2\mu(\mu - 1)\cos^{n\mu-2}(\vartheta z). \quad (17)$$

And so on for further derivatives.

We substitute Eq. (10) to (17) into the previously obtained reduced equation Eq. (8), balancing the terms of the cosine functions when Eq. (10) is used, or balancing the sine functions when Eq. (9) is used. Finally, we solve the resulting system of algebraic equations with the assistance of computerized symbolic computation to determine all possible values of the parameters λ, ϑ , and μ .

4. APPLICATION

In this section, we utilize the sine-cosine method to derive exact solutions for four distinct β -time fractional Boussinesq-like equations.

4.1 The First Time β – Fractional Boussinesq-like Equation

We consider The First Time β -Fractional Boussinesq-like Equation given by

$$D_{tt}^{2\beta} u - u_{xx} - (6u^2 u_x + u_{xxx})_x = 0. \quad (18)$$

By applying the following wave transformation:

$$u(x, t) = u(z),$$

$$\text{Where } z = \varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta, \quad (19)$$

Equation (18) reduces to ordinary differential equation:

$$\begin{aligned} \omega^2 u'' - \varpi^2 u'' - \varpi(6u^2 u_x + u_{xxx})' &= 0, \\ (\omega^2 - \varpi^2)u''(z) - \varpi(6\varpi u^2(z)u'(z) + \\ \varpi^3 u'''(z))' &= 0, \end{aligned} \quad (20)$$

Integrate once

$$(\omega^2 - \varpi^2)u'(z) - \varpi(6\varpi u^2(z)u'(z) + \varpi^3 u'''(z)) = 0, \quad (21)$$

Integrate again

$$\begin{aligned} (\omega^2 - \varpi^2)u(z) - \varpi \left(6\varpi \cdot \frac{u^3}{3} + \varpi^3 u''(z) \right) \\ = 0, \\ (\omega^2 - \varpi^2)u(z) - 2\varpi^2 u^3(z) - \varpi^4 u''(z) = 0. \end{aligned} \quad (22)$$

Eq. (22) is the reduced ordinary differential equation.

Assume that Eq. (22) has a solution in the form of

$$u(z) = \lambda \sin^\mu(\vartheta z), \quad (23)$$

$$u'(z) = \mu \lambda \vartheta \cos(\vartheta z) \cdot \sin^{\mu-1}(\vartheta z), \quad (24)$$

$$\begin{aligned} u''(z) &= \lambda \mu \vartheta (-\vartheta \mu \sin^\mu(\vartheta z) \\ &\quad + \vartheta (\mu - 1) \sin^{\mu-2}(\vartheta z)), \\ &= -\lambda \mu^2 \vartheta^2 \sin^\mu(\vartheta z) + \lambda \mu \vartheta^2 (\mu - 1) \sin^{\mu-2}(\vartheta z). \end{aligned} \quad (25)$$

For the cosine we have

$$u(z) = \lambda \cos^\mu(\vartheta z), \quad (26)$$

$$u'(z) = \lambda \mu \cos^{\mu-1}(\vartheta z) \cdot -\vartheta \sin^\mu(\vartheta z), \quad (27)$$

$$\begin{aligned} &= -\lambda \mu \vartheta \cos^{\mu-1}(\vartheta z) \sin^\mu(\vartheta z), \\ u''(z) &= -\lambda \mu^2 \vartheta^2 \cos^\mu(\vartheta z) + \lambda \vartheta^2 \mu (\mu - 1) \cos^{\mu-2}(\vartheta z). \end{aligned} \quad (28)$$

Substitute Eq. (23) and Eq. (25) in equation Eq. (22), we have

$$\begin{aligned} (\omega^2 - \varpi^2) \lambda \sin^\mu(\vartheta z) - 2 \varpi^2 \lambda^3 \sin^{3\mu}(\vartheta z) \\ - \varpi^4 \{-\lambda \mu^2 \vartheta^2 \sin^\mu(\vartheta z) \\ + \lambda \mu \vartheta^2 (\mu - 1) \sin^{\mu-2}(\vartheta z)\} = 0, \end{aligned} \quad (29)$$

$$\begin{aligned} (\omega^2 - \varpi^2) \lambda \sin^\mu(\vartheta z) - 2 \varpi^2 \lambda^3 \sin^{3\mu}(\vartheta z) \\ + \varpi^4 \lambda \mu^2 \vartheta^2 \sin^\mu(\vartheta z) \\ - \varpi^4 \lambda \mu \vartheta^2 (\mu - 1) \sin^{\mu-2}(\vartheta z) = 0. \end{aligned} \quad (30)$$

Eq. (30) is fulfilled if the following system of algebraic equations is satisfied:

$$\mu - 1 \neq 0,$$

$$\begin{aligned} 3\mu &= \mu - 2, \\ -2\varpi^2 \lambda^3 - \varpi^4 \lambda \mu \vartheta^2 (\mu - 1) &= 0, \\ (\omega^2 - \varpi^2) \lambda + \varpi^4 \lambda \mu^2 \vartheta^2 &= 0. \end{aligned}$$

Solving the system using Maple, we obtained

$$\mu = -1, \quad \lambda = \pm \frac{\sqrt{\varpi^2 - \omega^2}}{\varpi}, \quad \vartheta = \pm \frac{\sqrt{\varpi^2 - \omega^2}}{\varpi^2}. \quad (31)$$

Using the cosine method given by Eq. (26) will yield the same results.

Given Eq. (31), the solutions obtained from Eq. (23) and Eq. (19) are as follows:

$$u_{11}(x, t) = \frac{\sqrt{\varpi^2 - \omega^2}}{\varpi} \csc \left[\frac{\sqrt{\varpi^2 - \omega^2}}{\varpi^2} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right]. \quad (32)$$

And for cosine function, we have


$$u_{12}(x, t) = \frac{\sqrt{\varpi^2 - \omega^2}}{\varpi} \sec \left[\frac{\sqrt{\varpi^2 - \omega^2}}{\varpi^2} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right]. \quad (33)$$

Eq. (32) and Eq. (33) are valid only if $\varpi^2 - \omega^2 > 0$, so that $\omega^2 < \varpi^2$, $i = \sqrt{-1}$.


However, for $\omega^2 > \varpi^2$ we obtain the following solutions

$$u_{13}(x, t) = -\frac{\sqrt{\varpi^2 - \omega^2}}{\varpi} \operatorname{csch} \left[\frac{\sqrt{\varpi^2 - \omega^2}}{\varpi^2} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right], \quad (34)$$

$$u_{14}(x, t) = \frac{\sqrt{\varpi^2 - \omega^2}}{\varpi} \operatorname{sech} \left[\frac{\sqrt{\varpi^2 - \omega^2}}{\varpi^2} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right]. \quad (35)$$



(a) $\beta=0.5$






FIGURE 1. The graphical illustration for beta fractional derivative of solution $|u_{02}(x, t)|$ with $\varpi = 2, \omega = 1, t = 1.5, \mu = 1$. (a) $\beta = 0.5$, (b) $\beta = 0.75$, (c) $\beta = 0.95$, (d) Combined 2D graph for distinct values of β .

4.2 The Second Time β -Fractional Boussinesq-like Equation

In this section, we investigate the second β -time fractional Boussinesq-like equation:

$$D_{tt}^{2\beta} u - u_{xx} - (6u^2 u_x + D_{tt}^{2\beta} u_x)_x = 0. \quad (36)$$

We use the wave transformation

$$u(x, t) = u(z),$$

$$z = \varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta. \quad (37)$$

Eq. (35) reduces to

$$\omega^2 u'' - \varpi^2 u'' - \varpi (6u^2 u_x + u_{xtt})' = 0, \quad (38)$$

$$(\omega^2 - \varpi^2) u''(z) - \varpi (6\varpi u^2(z) u'(z) + \varpi \omega^2 u'''(z))' = 0, \quad (39)$$

Integrating Eq. (38) with respect to z once, we have

$$\begin{aligned} (\omega^2 - \varpi^2) u'(z) - \varpi (6\varpi u^2(z) u'(z) + \varpi \omega^2 u'''(z)) &= 0, \\ (\omega^2 - \varpi^2) u'(z) - 6\varpi^2 u^2(z) u'(z) - \varpi^2 \omega^2 u'''(z) &= 0, \end{aligned} \quad (40)$$

Integrating Eq. (40) with respect to z again, we obtain

$$(\omega^2 - \varpi^2) u(z) - 2\varpi^2 u^3(z) - \varpi^2 \omega^2 u''(z) = 0. \quad (41)$$

Now, Eq. (41) is the reduced ordinary

Solving system Eq. (44), we obtain

$$\begin{aligned} \mu &= -1, & \lambda &= \pm \frac{\sqrt{\omega^2 - \varpi^2}}{\varpi}, \\ \vartheta &= \pm \frac{\sqrt{\varpi^2 - \omega^2}}{\varpi \omega}, \end{aligned}$$

Similar results are also achieved when employing the cosine function.

As a result, the following solutions were derived:




FIGURE 2. The graphical illustration for beta fractional derivative of solution $|u_{21}(x, t)|$ with $\varpi = 2, \omega = -3, t = 1.5, \mu = -1$. (a) $\beta = 0.5$, (b) $\beta = 0.75$, (c) $\beta = 0.95$, (d) Combined 2Dgraph for distinct values of β .

$$u_{21}(z) = \frac{\sqrt{\omega^2 - \varpi^2}}{\varpi} \csc \left[\frac{\sqrt{\omega^2 - \varpi^2}}{\varpi \omega} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right], \quad \omega^2 < \varpi^2, \quad (45)$$

$$u_{22}(z) = \frac{\sqrt{\omega^2 - \varpi^2}}{\varpi} \sec \left[\frac{\sqrt{\omega^2 - \varpi^2}}{\varpi \omega} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right], \quad \omega^2 < \varpi^2. \quad (46)$$

For $\omega^2 > \varpi^2$ we also obtained the following solutions:

$$u_{23}(z) = -i \frac{\sqrt{\omega^2 - \varpi^2}}{\varpi} \operatorname{csch} \left[\frac{\sqrt{\omega^2 - \varpi^2}}{\varpi \omega} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right], \quad (47)$$

$$u_{24}(z) = \frac{\sqrt{\omega^2 - \varpi^2}}{\varpi} \operatorname{sech} \left[\frac{\sqrt{\omega^2 - \varpi^2}}{\varpi \omega} \left(\varpi x + \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right]$$

$$\varpi^3 \omega \lambda \mu^2 \vartheta^2 \sin^\mu(\vartheta z) + \varpi^3 \omega \lambda \mu \vartheta^2 (\mu - 1) \sin^{\mu-2}(\vartheta z) = 0. \quad (53)$$

By equating the exponents and coefficients of each pair of sine functions to zero, we derive the following system of algebraic equations:

$$\begin{aligned} \mu - 1 &\neq 0, \\ 3\mu &= \mu - 2, \\ (\omega^2 + \varpi\omega)\lambda - \varpi^3 \omega \lambda \mu^2 \vartheta^2 &= 0, \end{aligned} \quad (54)$$

$$-2\varpi^2 \lambda^3 + \varpi^3 \omega \lambda \mu \vartheta^2 (\mu - 1) = 0.$$

Solving the system Eq. (54) yields

$$\mu = -1, \quad \lambda = \pm \frac{\sqrt{\omega^2 + \omega\varpi}}{\varpi}, \quad \vartheta = \pm \frac{\sqrt{\varpi(\omega + \varpi)}}{\varpi^2}. \quad (55)$$

Consequently, for $\varpi\omega + \varpi^2 > 0$, we obtained the following solutions

$$u_{31}(z) = \frac{\sqrt{\omega^2 + \omega\varpi}}{\varpi} \csc \left[\frac{\sqrt{\varpi(\omega + \varpi)}}{\varpi^2} \left(\varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right], \quad (56)$$

and

$$u_{32}(z) = \frac{\sqrt{\omega^2 + \omega\varpi}}{\varpi} \sec \left[\frac{\sqrt{\varpi(\omega + \varpi)}}{\varpi^2} \left(\varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right]. \quad (57)$$

However, for $\varpi\omega + \varpi^2 < 0$, gives the following solutions

$$u_{33}(z) = -i \frac{\sqrt{-\omega^2 - \omega\varpi}}{\varpi} \operatorname{csch} \left[\frac{\sqrt{-\varpi(\omega + \varpi)}}{\varpi^2} \left(\varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right], \quad (58)$$

and

$$u_{34}(z) = \frac{\sqrt{-\omega^2 - \omega\varpi}}{\varpi} \operatorname{sech} \left[\frac{\sqrt{-\varpi(\omega + \varpi)}}{\varpi^2} \left(\varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right]. \quad (59)$$

4.3 The Third Time β –Fractional Boussinesq-like Equation

In this section we study the third time beta-fractional Boussinesq-like equation:

$$D_{tt}^{2\beta} u - u_{xt}^\beta - (6u^2 u_x + D_t^\beta u_{xx})_x = 0. \quad (49)$$

Using the travelling wave transformation:

$$u(x, t) = u(z),$$

$$z = \varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta.$$

Substituting these in Eq. (49) gives

$$(\omega^2 + \varpi\omega)u''(z) - \varpi(6u^2(z)u'(z) - \varpi^2 \omega u'''(z))' = 0, \quad (50)$$

Integrating twice and setting the constants of integration to zero, we obtain:

$$(\omega^2 + \varpi\omega)u(z) - 2\varpi^2 u^3(z) + \varpi^3 \omega u''(z) = 0. \quad (51)$$

Assuming that Eq. (51) has a solution in the form:

$$u(z) = \lambda \sin^\mu(\vartheta z),$$

Where ω and ϖ are constants, substituting Eq. (23) and Eq. (25) into Eq. (51) yields:

$$(\omega^2 + \varpi\omega)\lambda \sin^\mu(\vartheta z) - 2\varpi^2 (\lambda \sin^\mu(\vartheta z))^3 + \varpi^3 \omega (-\lambda \mu^2 \vartheta^2 \sin^\mu(\vartheta z) + \lambda \mu \vartheta^2 (\mu - 1) \sin^{\mu-2}(\vartheta z)) = 0, \quad (52)$$

$$(\omega^2 + \varpi\omega)\lambda \sin^\mu(\vartheta z) - 2\varpi^2 \lambda^3 \sin^{3\mu}(\vartheta z) -$$




FIGURE 3. The graphical illustration for beta fractional derivative of solution $|u_{31}(x, t)|$ with $\varpi = 2, \omega = 1, t = 1.5, \mu = -1$. (a) $\beta = 0.5$, (b) $\beta = 0.75$, (c) $\beta = 0.95$, (d) Combined 2D graph for distinct values of β .

4.4 The Fourth Time β –Fractional Boussinesq- like Equation

In this section we study the fourth time beta-fractional Boussinesq like equation:

$$D_{tt}^{2\beta} u - (6u^2 u_x + u_{xxx})_x = 0. \quad (60)$$

Using the travelling wave transformation:

$$u(x, t) = u(z),$$

$$z = \varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta.$$

Substituting these in Eq. (60) becomes

$$\omega^2 u''(z) - \varpi (6u^2 u_x + u_{xxx})' = 0, \quad (61)$$

$$\omega^2 u''(z) - \varpi (6u^2 \varpi u'(z) + \varpi^3 u'''(z))' = 0, \quad (62)$$

Integrating Eq. (62) once, we have

$$\omega^2 u'(z) - \varpi (6\omega u^2(z) u'(z) + \varpi^3 u'''(z)) = 0, \quad (63)$$

$$\omega^2 u'(z) - 6\varpi^2 u^2(z) u'(z) - \varpi^4 u'''(z) = 0, \quad (64)$$

Integrating Eq. (64) again gives

$$\omega^2 u(z) - 2\varpi^2 u^3(z) - \varpi^4 u''(z) = 0. \quad (65)$$

Where ϖ and ω are constants. Substituting Eq. (23) and Eq. (25) into Eq. (65) gives

$$\omega^2 \lambda \sin^\mu(\vartheta z) - 2\varpi^2 \lambda^3 \sin^{3\mu}(\vartheta z) - \varpi^4 (-\lambda \mu^2 \vartheta^2 \sin^\mu(\vartheta z) + \lambda \mu \vartheta^2 (\mu - 1) \sin^{\mu-2}(\vartheta z)) = 0, \quad (66)$$

$$\omega^2 \lambda \sin^\mu(\vartheta z) - 2\varpi^2 \lambda^3 \sin^{3\mu}(\vartheta z) + \varpi^4 \lambda \mu^2 \vartheta^2 \sin^\mu(\vartheta z) - \varpi^4 \lambda \mu \vartheta^2 (\mu - 1) \sin^{\mu-2}(\vartheta z) = 0. \quad (67)$$

(67)

By setting the exponents and coefficients of each pair of sine functions to zero, we derive the following system of algebraic equations:




FIGURE 4. (a)




FIGURE 4. (b)




FIGURE 4. (c)




FIGURE 4. (d)

FIGURE 4. The graphical illustration for beta fractional derivative of solution $|u_{41}(x,t)|$ with $\varpi = 5$, $\omega = 3$, $t = 1.5$, $\mu = -1$. (a) $\beta = 0.5$, (b) $\beta = 0.75$, (c) $\beta = 0.95$, (d) Combined 2D graph for distinct values of β .

$$\begin{aligned} \mu - 1 &\neq 0, \\ 3\mu &= \mu - 2, \\ \omega^2\lambda + \varpi^4\lambda\mu^2\vartheta^2 &= 0, \\ (68) \quad -2\varpi^2\lambda^3 - \varpi^4\lambda\mu\vartheta^2(\mu - 1) &= 0. \end{aligned}$$

Upon solving the system of equations Eq. (68), we obtain:

$$\mu = -1, \quad \lambda = \pm \frac{\omega}{\varpi}, \quad \vartheta = \pm \frac{\omega i}{\varpi^2}. \quad (69)$$

Consequently, we obtained the following solutions

$$u_{41}(z) = \frac{\omega}{\varpi} \csc \left[\frac{\omega i}{\varpi^2} \left(\varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right)^\beta \right) \right], \quad (70)$$

and

$$u_{42} = \frac{\omega}{\varpi} \sec \left[\frac{\omega i}{\varpi^2} \left(\varpi x - \frac{\omega}{\beta} \left(t + \frac{1}{\Gamma(\beta)} \right) \right) \right]. \quad (71)$$

5 PHYSICAL EXPLANATION

In this section, we generated various types of exact solutions for Equations (1)-(4) using the Sine-cosine approach. These solutions are in the form of trigonometric and hyperbolic functions.

Fig 1. (a)-(c) Depicts modulus of solutions for $|u_{11}(x,t)|$, Eq. (32) with various parameter values, $\varpi = 2$, $\omega = 1$, $\mu = -1$, $\lambda = \frac{\sqrt{3}}{2}$, $\vartheta = \frac{\sqrt{6}}{4}$, as well as the beta fractional order values, $\beta = 0.5$ for (a), $\beta = 0.75$ for (b) and $\beta = 0.95$ for (c). We observed multiple solitons with varying heights and positions. (d) Shows the effect of beta fractional order values in combined 2D wave profile with $\beta = 0.5$, 0.75 , 0.95 and $t = 1.5$. We observed the wave propagates in the x-direction for increasing values of

β . For specific values of the parameters $\varpi = 2$, $\omega = -3$, $\mu = -1$, $\lambda = \frac{\sqrt{5}}{2}$, $\vartheta = -\frac{\sqrt{5}}{6}I$, the 3D wave profile of the solution $|u_{2,1}(x,t)|$ which depicts periodic solitary wave corresponding to Fig 2.(a)-(c) with different beta values $\beta = 0.5$ for (a), $\beta = 0.75$ for (b) and $\beta = 0.95$ for (c). Moreover, the combined 2D graph is shown in Fig. 2(d) with distinct beta values, $\beta = 0.5, 0.75, 0.95$ and $t = 1.5$.

Fig. 3(a)-(c) represents the periodic wave structure of the solution $|u_{3,1}(x,t)|$ for the parameter values $\varpi = 2$, $\omega = 1$, $\mu = -1$, $\lambda = \frac{\sqrt{3}}{2}$, $\vartheta = \frac{\sqrt{6}}{4}$, with $\beta = 0.5$ for (a), $\beta = 0.75$ for (b) and $\beta = 0.95$ for (c) and the 2D graph is plotted in Fig. 3(d) with beta values $\beta = 0.5, 0.75, 0.95$ and $t = 1.5$, which shows the effect of beta fractional order. Fig. 4(a)-(c) illustrates the 3D wave structure of the solution $|u_{4,1}(x,t)|$ representing singular soliton solution with parameter values $\varpi = 5$, $\omega = 3$, $\mu = -1$, $\lambda = \frac{3}{5}$, $\vartheta = 3/25$, and for specific values of the beta: $\beta = 0.5$ for (a), $\beta = 0.75$ for (b) and $\beta = 0.95$ for (c). The 2D combined wave profiles are shown in Fig. 4(d) for $\beta = 0.5, 0.75, 0.95$ and $t = 1.5$ to show the effect of beta fractional order. From the above discussion, it is obvious that the sine-cosine approach can provide different types of wave structures to fractional Boussinesq-like equations with beta derivative for various values of parameters.

The obtained solutions derived through this method play a crucial role in understanding the structure and dynamic behavior of the problem at hand. The method proposed in this work is straightforward, dependable, and effective, with the potential to be extended for studying and solving numerous other nonlinear evolution equations across various scientific and engineering disciplines.

CONCLUSION

In this paper, we employed the sine-cosine method to obtain travelling wave solutions for four distinct Beta fractional Boussinesq-like equations. The obtained results include singular

traveling wave soliton solution, periodic soliton solutions and multiple soliton solution which involve trigonometric functions, hyperbolic functions and complex solutions as well. We examined all constraints that guarantee the existence of these new exact solutions. The results obtained from this study are anticipated to be highly beneficial in numerous areas of mathematical physics, costal engineering and applied mathematics, including fluid dynamics, nonlinear optics, plasma physics, and other related fields.

ACKNOWLEDGEMENT

The author would like to express his gratitude to the editor of the journal and the anonymous reviewer for taking their precious time in providing detailed comments and suggestions that really helped to improve the quality of the paper.

REFERENCES

- Ahmad, J., Rani, S., Turki, N. B. and Shah, N. A. (2023). *Novel resonant multi-soliton solutions of time fractional coupled nonlinear Schrodinger equation in optical fiber via an analytical method*. Results Phys52. 106761.
- Ahmad, J., Akram, S. U., Turki, N. B. and Shah, N.A. (2023). *Description of soliton and lump solutions to M-truncated stochastic Biswas-Arshed model I optical communication*. Results Phys. 51, 10679.
- Akram, S., Ahmad, J., Sarwar, S. and Ali, A. (2023). *Dynamic of soliton solution in optical fiber modelled by perturbed nonlinear Schrodinger equation and stability analysis*. Opt. Quantum. Electronics. 55 (5) 450.
- Ali, A., Ahmad, J. and Javed, S. (2023). *Exploring the dynamic nature of soliton solutions to the coupled nonlinear Schrodinger equation with their sensitivity analysis*. Opt. Quantum Electron. 55(9), 810.
- Ali, A., Ahmad, J. and Javed, S. (2023). *Investigating the dynamics of soliton solutions to the fractional coupled nonlinear Schrodinger model with their bifurcation and stability analysis*. Opt. Quantum Electron, 55(9), 829.
- Atangana, A. and Baleanu, D. (2016). *New fractional*

- derivatives with nonlocal and non-singular kernel. Theory and application to heat transfer model. Therm. Sci. 20 (2), 763-769.
- Atangana A, Baleanu D, and Alsaedi A. (2016). *Analysis of time-fractional Hunter-Saxton equation: a model of neumatic liquid crystal*. Open Phys. 14(1), 145-149 doi: 10.1515/phys-2016-0010.
- Darvishi, M. T., Naja, M. and Wazwaz, A. M. (2017). *Soliton solutions for Boussinesq-like equations with spatio-temporal dispersion*, Ocean Eng. 130, 228-240.
- Darvishi, M. T., Naja, M. and Wazwaz, A. M. (2018). *Traveling wave solutions for Boussinesq-like Equations with Spatial and Spatial-Temporal Dispersion*. Romanian Reports in Physics, 70, 108.
- Eslami, M. and Mirzazadeh, M. (2014). *First integral method to look for exact solutions of a variety of Boussinesq-like equations*, Ocean Eng. 83, 133-137.
- Fiza, B., Muhammad, S. S., Ulviye, D., Hadi, R., Khalid, M. K., Saleh, A. and Hijaz, A. (2024). *Studying the impact of M-fractional and beta derivatives on the nonlinear fractional mode*. Optical and Quantum Electronics 56:164.
- Ghanbari, B. (2019). *Abundant soliton solution for the Hirota- Maccari equation via the generalized exponential rational function method*. Mod. Phys. Lett. B. 33 (09), 1950106.
- Ghanbari, B. and Baleanu, D. (2020). *New optical solutions of the fractional Gerdjikov-Ivanov equation with conformable derivative*. Front. Phys. 8: 167.
- Jumarie, G. (2009). *Table of some basic fractional calculus formulae derived from a modified Riemann-Liouville derivative for non-differentiable functions*. Appl. Math. Lett. 22 (3), 378-385.
- Khalil, R., Al Horani, M., Yousef, A. and Sababheh, M. (2014). *A new definition of fractional derivative*. J. Comput. Appl. Math. 264, 65-70.
- Khater, M. and Ghanbari, B. (2021). *On the solitary wave solution and physical characterization of gas diffusion in a homogenous medium via some efficient techniques*. Eur. Phys. J. 134 (4), 1-28.
- Khatun, M. A., Arefin, M. A., Uddin, M. H., Inc. M. and Akbar, M. A. (2022). *An analytical approach to the solution of fractional coupled modified equal width and fractional coupled Burgers equations*. J. Ocean Eng. Sci.
- Osman, M. S. (2019). *On multi-soliton solutions for the (2+1) - dimensional breaking soliton equation with variable coefficients in a graded-index waveguide*, Comput. Math. Appl. 75(1), 1-6.
- Sadiya, U., Arefin, M. A. and Uddin, M. H. (2022). *Consistence travelling wave solutions to the nonlinear time fractional Klein- Gordon and Sine-Gordon equations thorough extended tanh function approach*, J. Taibah Univ. Sci. 16 (1), 594-607.
- Sousa, J. V. D. C. and de Oliveira, E. C. (2018). *A new truncated M-fractional derivative type unifying some fractional derivative types with classical properties*. Int. J. Anal. Appl. 16 (1), 83-96.
- Uddin, M. H., Zaman, U. H. M. and Arefin, M. A. (2022). *Nonlinear dispersive wave propagation pattern in optical fiber system*, Chaos solitons Fractals 164, 112596.
- Wazwaz, A. M. (2004). Math. Comput. Model. 40 (5-6), 499-508.
- Wazwaz, A. M. (2012). *Solitons and singular solitons for a variety of Boussinesq-like equations*, Ocean. 53, 1-5.
- Yepez-Martinez, H., Gomez-Aguila, J. F. and Baleanu, D. (2018). *Beta derivative and sub-equation method applied to the optical soliton in medium with parabolic law nonlinearity and higher order dispersion*. Optik 155, 357-365.

Yiasir Arafat, S. M., Islam, R. S. M., Rahman, M. M. and Saklayen, M. A. (2023). *On nonlinear Optical Solitons of Fractional Biswas-Arshed Model with Beta Derivative*. Results I Physics 48 106426.

Yusuf, A., Inc, M. and Baleanu, D. (2019). *Optical soliton with M-truncated and beta derivatives in nonlinear optics*. Front. Phys. 7, 126.

Zaman, U. H. M., Arefin, M. A., Akbar, M. A. and Uddin, M. H. (2022a). *Analyzing numerous travelling wave bahaviour to the fractional – order nonlinear Phi-4 and Allen –Cahn equations throughout a novel technique*. Results Phys. 37, 105486.

Zaman U. H. M., Arefin, M. A. and Akbar, M. A. (2022b). *Analytical behavior of soliton solutions to the couple type fractional order nonlinear evolution equations utilizing a novel technique*. Alex. Eng. J. 61(12), 11947-11958.